


AMENDMENTS TO THE CLAIMS

1. (Original) A fungicidal mixture, comprising


- 1) the triazolopyrimidine derivative of the formula I,



I

and

- 2) fenpropimorph of the formula II,



II

in a synergistically effective amount.

2. (Original) A fungicidal mixture, comprising the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.

3. (Original) A fungicidal composition, comprising a liquid or solid carrier and a mixture as claimed in claim 1 or 2.

4. (Currently Amended) A method for controlling harmful fungi which are rice pathogens, which comprises treating the fungi, their habitat or the plants, the soil or the seeds to be protected

against fungal attack with an effective amount of the compound I and the compound II as set forth in claim 1 ~~or of the composition as claimed in claim 3.~~

5. (Currently Amended) A method ~~as claimed in claim 4~~ for controlling harmful fungi which are rice pathogens, which comprises treating the fungi, their habitat or the plants, the soil or the seeds to be protected against fungal attack with an effective amount of the compound I and the compound II as set forth in claim 1, wherein the compounds I and II ~~as set forth in claim 1~~ are applied simultaneously, that is jointly or separately, or in succession.

6. (Currently Amended) A method as claimed in claim 4, wherein the mixture as claimed in claim 1 or 2 ~~or the composition as claimed in claim 3~~ is applied in an amount of from 5 g/ha to 2000 g/ha.

7. (Currently Amended) A method as claimed in claim 4 ~~or 5~~, wherein the mixture as claimed in claim 1 or 2 ~~or the composition as claimed in claim 3~~ is applied in an amount of from 0.001 to 1 g/kg of seed.

8. (Currently Amended) A method as claimed in ~~any of claims 4 to 7~~ claim 4, wherein the harmful fungus *Corticium sasakii* is controlled.

9. (Original) Seed, comprising the mixture as claimed in claim 1 or 2 in an amount of from 0.001 to 1 g/kg.

10. (Original) The use of the compound I and the compound II as set forth in claim 1 for preparing a composition suitable for controlling harmful fungi.

11. (New) A method as claimed in claim 5, wherein the mixture as claimed in claim 1 or 2 is applied in an amount of from 0.001 to 1 g/kg of seed.